19. (Amended) A pharmaceutical composition comprising a compound as claimed in claim 1 and a pharmaceutically acceptable carrier.

20. (Amended) A method of treating or preventing an integrin-mediated disease or condition, comprising administering to a mammal an effective amount of a compound as claimed in claim 1.

21. (Amended) A method of inhibiting angiogenesis and/or for treating or preventing cancer, osteolytic diseases and ophthalmic disorders, comprising administering to a mammal an effective amount of a compound of the general formula (1)



(1)

where

R^1 is hydrogen, a substituted or unsubstituted alkyl or cycloalkyl residue, a substituted or unsubstituted aryl residue or a saturated or unsaturated, optionally substituted heterocyclic residue;

R^2 is hydrogen, a substituted or unsubstituted alkyl or cycloalkyl residue, a substituted or unsubstituted aryl residue, a saturated or unsaturated, optionally substituted heterocyclic residue, an optionally substituted alkenyl residue, an optionally substituted alkinyl residue, $-NR^{2'}SO_2R^{2''}$, $-NR^{2'}COOR^{2'}$, $-NR^{2'}COR^{2'}$, $-NR^{2'}CONR^{2'}_2$ or $-NR^{2'}CSNR^{2'}_2$;

$R^{2'}$ is hydrogen, a substituted or unsubstituted alkyl or cycloalkyl residue, a substituted or unsubstituted aryl residue or a saturated or unsaturated, optionally substituted heterocyclic residue;

$R^{2'}$ is hydrogen, a substituted or unsubstituted alkyl or cycloalkyl residue, a substituted or unsubstituted aryl residue or a saturated or unsaturated, optionally substituted heterocyclic residue;

$R^{2''}$ is a substituted or unsubstituted alkyl, alkenyl or cycloalkyl residue, a substituted or unsubstituted aryl residue or a saturated or unsaturated, optionally substituted heterocyclic residue;

U is a direct bond or a substituted or unsubstituted alkylene group;

V is a substituted or unsubstituted alkylene group, $-NR^{2'}CO-$ or $-NR^{2'}SO_2-$;

A and B each independently of one another a 1,3- or 1,4-bridging phenylene group or a 2,4- or 2,5-bridging thienylene group each of which may optionally have additional substituents,

W is a direct bond or a substituted or unsubstituted alkylene group;

C is a direct bond or  or

R^3 is hydrogen, a substituted or unsubstituted alkyl or cycloalkyl residue, a substituted or unsubstituted aryl residue, a saturated or unsaturated, optionally substituted heterocyclic residue, an alkylamine residue, an alkylamide residue or is connected to one of R^4, Y, R^5 or R^6, if present, with formation of an optionally substituted heterocyclic ring system which includes the nitrogen atom to which R^3 is bonded, and can be saturated or unsaturated and/or can contain further heteroatoms;

R^4 is hydrogen, a substituted or unsubstituted alkyl or cycloalkyl residue, a substituted or unsubstituted aryl residue, a saturated or unsaturated, optionally substituted heterocyclic residue, an alkylamine residue, an alkylamide residue

X is $CHNO_2$, CHCN. O, N or S;

Y is a direct bond or an optionally substituted alkylene or alkine group;

R^5 is absent, or is hydrogen, a substituted or unsubstituted alkyl or cycloalkyl residue, $-NO_2$, -CN, $-COR^{5'}$, $-COOR^{5'}$, or is connected to one of R^3, Y, R^4 or R^6, if present, with formation of an optionally substituted carbocyclic or heterocyclic ring system which includes X and can be saturated or unsaturated and/or can contain further heteroatoms;

$R^{5'}$ is hydrogen, a substituted or unsubstituted alkyl or cycloalkyl residue, a substituted or unsubstituted aryl residue or a saturated or unsaturated, optionally substituted heterocyclic residue which can be saturated or unsaturated and/or can contain further heteroatoms;

R^6 is hydrogen, a substituted or unsubstituted alkyl or cycloalkyl residue, a substituted or unsubstituted aryl or aroyl residue, a saturated or unsaturated, optionally substituted heterocyclic residue, an alkylamine residue, an alkylamide residue or is connected to one of R^3, R^4, Y or R^5, if present, with formation of an optionally substituted heterocyclic ring system which includes the nitrogen atom to which R^6 is bonded and can be saturated or unsaturated and/or can contain further heteroatoms;

and their physiologically acceptable salts and stereoisomers.

22. canceled

New Claim for Attorney Docket No. Le A 33 324(Nat'l Stage of PCT/EP99/09843)

23. (New) The method of claim 21, wherein said osteolytic disease is selected from the group consisting of osteoporosis, arteriosclerosis, restenosis and rheumatoid arthritis.